

Role of transporters in the distribution of platinum-based drugs

Saliha Harrach¹ and Giuliano Ciarimboli^{1,2*}

¹ Experimental Nephrology, Medical Clinic D, University of Münster, University Hospital Münster, Münster, Germany,

² Interdisciplinary Center for Clinical Research (IZKF), University of Münster, University Hospital Münster, Münster, Germany

OPEN ACCESS

Edited by:

Harry P. De Koning,
University of Glasgow, UK

Reviewed by:

Stanislav Yanev,
Bulgarian Academy of Sciences,
Bulgaria
Andrei Adrian Tica,
University of Medicine Craiova
Romania, Romania

*Correspondence:

Giuliano Ciarimboli,
Experimental Nephrology, Medical
Clinic D, Münster University Hospital,
Albert-Schweitzer-Campus 1–A14,
48149 Münster, Germany
gciari@uni-muenster.de

Specialty section:

This article was submitted to
Drug Metabolism and Transport,
a section of the journal
Frontiers in Pharmacology

Received: 27 February 2015

Paper pending published:
30 March 2015

Accepted: 02 April 2015

Published: 24 April 2015

Citation:

Harrach S and Ciarimboli G (2015)
Role of transporters in the distribution
of platinum-based drugs.
Front. Pharmacol. 6:85.
doi: 10.3389/fphar.2015.00085

Platinum derivatives used as chemotherapeutic drugs such as cisplatin and oxaliplatin have a potent antitumor activity. However, severe side effects such as nephro-, oto-, and neurotoxicity are associated with their use. Effects and side effects of platinum-based drugs are in part caused by their transporter-mediated uptake in target and non target cells. In this mini review, the transport systems involved in cellular handling of platinum derivatives are illustrated, focusing on transporters for cisplatin. The copper transporter 1 seems to be of particular importance for cisplatin uptake in tumor cells, while the organic cation transporter (OCT) 2, due to its specific organ distribution, may play a major role in the development of undesired cisplatin side effects. In polarized cells, e.g., in renal proximal tubule cells, apically expressed transporters, such as multidrug and toxin extrusion protein 1, mediate secretion of cisplatin and in this way contribute to the control of its toxic effects. Specific inhibition of cisplatin uptake transporters such as the OCTs may be an attractive therapeutic option to reduce its toxicity, without impairing its antitumor efficacy.

Keywords: cisplatin, oxaliplatin, transporters, side effects, uptake

In order to reach sufficiently high concentrations, intracellular active drugs must cross the plasma membrane to reach their targets. This process, especially in the case of hydrophilic drugs, is mediated by specialized plasma membrane proteins called transporters. It is well known that efflux transporters are involved in the development of drug resistance (Renes et al., 2000). Only recently, it has become evident that uptake transporters not only are able to mediate drug effects but also their side effects (Ciarimboli, 2011). In this mini-review, we will focus on transporter systems for platinum-based drugs.

Cisplatin was the first platinum-based drug that revolutionized the treatment of neoplastic diseases. For example, before the introduction of cisplatin as chemotherapeutic agent, testicular cancer was associated with a survival rate of only 5%. Today, treatment of this cancer with a combination of new surgical techniques and cisplatin chemotherapy allows to achieve a cure rate of over 90% (Einhorn, 2002). Currently, cisplatin is widely used for the therapy of solid tumors. However, its use is limited by severe side effects such as nephro- and oto-toxicity and peripheral neurotoxicity (Rabik and Dolan, 2007). Therefore, there is a need to put an effort in developing less toxic platinum derivatives. The US Food and Drug Administration has approved carboplatin and oxaliplatin as chemotherapeutic agents. Nevertheless, because of its superior efficacy, cisplatin remains the first-line treatment for several solid cancers.

Treatment with cisplatin causes an acute and/or chronic nephrotoxicity in about one-third of patients even under co-treatment with diuretics and pre-hydration (Pabla and Dong, 2008). The clinical manifestations of cisplatin nephrotoxicity are a decreased glomerular filtration rate,

increased serum creatinine level, reduced serum magnesium and potassium levels (Pabla and Dong, 2008), and the development of proteinuria of tubular and glomerular origin (Daugaard, 1990). There are striking interindividual differences in susceptibility to nephrotoxicity (Goren et al., 1986).

Animal studies showed that cisplatin damages mainly the proximal tubules (Dobyan et al., 1980), where the S3 segment is highly sensitive to cisplatin toxicity and undergoes extensive necrosis *in vivo* (Price et al., 2004). In contrast, carboplatin treatment rarely results in nephrotoxicity (Bregman and Williams, 1986; Yasumasu et al., 1992; Rabik and Dolan, 2007) and nephrotoxicity has never been observed with oxaliplatin, allowing its administration without hydration (Cassidy and Misset, 2002).

Tumor therapy with cisplatin causes a hearing loss, which is an unresolved clinical problem especially in pediatric patients (Skinner, 2004). By early onset, cisplatin therapy may lead to a delayed speech development with serious consequences for social and educational development (Skinner, 2004). Ototoxicity appears with an incidence between 23 and 50% in adults and greater than 50% in children (Rabik and Dolan, 2007). It manifests clinically with bilateral symmetrical high-frequency sensorineural hearing loss, ear pain or tinnitus (Reddel et al., 1982). In the cochlea cisplatin accumulates in the hair cells of the basal turn of Corti organ (where high-frequency sounds are processed), in the spiral ganglion cells, and in the lateral wall tissues (spiral ligament and stria vascularis) (Rybak and Ramkumar, 2007). Upon cisplatin accumulation, these cells undergo apoptosis (Rybak and Ramkumar, 2007). Carboplatin and oxaliplatin are much less ototoxic than cisplatin (Musial-Bright et al., 2011). Interestingly, intravenous application of oxaliplatin (16.6 mg/kg) in guinea pigs resulted in cochlear drug concentrations lower than those measured upon treatment with equimolar cisplatin (12.5 mg/kg), suggesting that lower cochlear oxaliplatin accumulation is the reason for its lower ototoxicity compared with cisplatin (Hellberg et al., 2009).

Most patients treated with cisplatin develop a dose-related large fiber sensory neuropathy, caused by cisplatin accumulation in the dorsal root ganglia (DRG) (Meijer et al., 1999; Albers et al., 2011). The symptoms of peripheral neurotoxicity may appear already 1 month after initiating treatment and include paresthesia, ataxia, numbness, reduced vibration, and joint position sensations and diminished or absent muscle stretch reflexes (Roelofs et al., 1984; Thompson et al., 1984; Cano et al., 1998; Albers et al., 2011). Cisplatin toxicity seems to affect large diameter neurons and proprioceptive sensory modalities (reviewed in Screnci and McKeage, 1999), where platinum accumulation within DRG leads to atrophy or loss of peripheral sensory neurons (Liu et al., 2009). Carboplatin is less neurotoxic than cisplatin. Neurotoxicity appears in only 4–6% of patients after administration of high-dose carboplatin therapy (Amptoulach and Tsavaris, 2011). Conversely, antitumor treatment with oxaliplatin causes significant acute and chronic peripheral sensory neuropathy (Amptoulach and Tsavaris, 2011). Acute neurotoxic effects develop in 85–95% of patients treated with oxaliplatin and may result from the impairment of voltage-gated sodium channels, reduction in intracellular calcium and of growth-associated protein-43

expression, or induction of oxidative stress (Adelsberger et al., 2000; Gamelin et al., 2002; Carozzi et al., 2010; Amptoulach and Tsavaris, 2011; Takeshita et al., 2011). The symptoms of oxaliplatin neurotoxicity include paresthesia in the extremities and the perioral region, and dysfunction of fine sensory-motor coordination (Cavaletti and Zanna, 2002; Hartmann and Lipp, 2003; Amptoulach and Tsavaris, 2011). These symptoms are exacerbated by cold exposure (Amptoulach and Tsavaris, 2011). It is evident that oxaliplatin neurotoxicity impairs quality of daily life.

The antitumor effects of platinum derivatives seem to derived from the formation of intra- and inter-strand crosslinks with DNA (Wang and Lippard, 2005).

Focusing on cisplatin, the molecular species involved in this process seems to be aquaions, which are supposed to be formed upon cytosolic cisplatin hydrolysis driven by the lower intracellular Cl^- concentration. The cisplatin aquaions are strongly electrophilic and bind DNA and sulfhydryl groups of proteins.

Since cisplatin is a hydrophilic drug (Houjou et al., 1996), it cannot easily cross the plasma membrane. In the kidneys cisplatin is free filtered in the glomerulus and also secreted into the urine (Jacobs et al., 1984). These observations suggested that the movement of cisplatin through the plasma membrane is mediated by transport proteins.

Copper Transporter 1 (Ctr1, Solute Carrier 31A1-SLC31A1)

Copper transporter 1 (Ctr1, Solute Carrier 31A1-SLC31A1) is a membrane protein that plays a significant role in the cellular cisplatin uptake (Holzer et al., 2004, 2006; Safaei and Howell, 2005; Larson et al., 2009). Down-regulation of Ctr1 extensively reduced cisplatin uptake in yeast and in mouse embryonic fibroblasts (Ishida et al., 2002; Lin et al., 2002). The natural substrate of Ctr1 is monovalent copper (Cu^+) (Ohrvik and Thiele, 2014). Cu^+ uptake by Ctr1 triggers transporter internalization. However, whether this phenomenon also happens upon cisplatin transport is debated (Sinani et al., 2007; Tsai et al., 2014). As observed for Cu^+ , cisplatin binds to Methionine-rich motifs of the extracellular domain of Ctr1 (Ohrvik and Thiele, 2014). Ctr1 carries out vital physiological function supplying the cell with copper, which is an essential cellular nutrient used in a broad range of enzymatic reactions. Because of its important biological role, Ctr1 is almost ubiquitously expressed and perhaps may not be the decisive transporter for specific cisplatin toxicities. Since several cell lines from human tumor samples express Ctr1-mRNA (Ciarimboli et al., 2010), this transporter could represent the uptake route of cisplatin in cancer cells. Indeed, high expression levels of Ctr1 have been associated with cisplatin therapeutic success (Lee et al., 2011; Liang et al., 2012) whereas Ctr1 mutations are associated with cisplatin resistance (Xu et al., 2012). Ctr1 has been also associated with the cellular transport of carboplatin and oxaliplatin. However, cells with a genetic deletion of Ctr1 show an uptake of oxaliplatin at concentrations higher than $2 \mu\text{M}$, suggesting the presence of another low-affinity oxaliplatin transport system (Holzer et al., 2006).

Copper Transporter 2 (Ctr2, SLC31A2)

Copper transporter 2 (Ctr2, SLC31A2) is a copper transport protein with substantial structural homology to Ctr1. Ctr2 is mainly expressed in late endosomes and lysosomes (Blair et al., 2009), where it probably mediates the efflux of copper under conditions of low environmental copper concentration (van den Berghe et al., 2007). A similar function of Ctr2 was proposed for cisplatin (Blair et al., 2011). Studies in Ctr2-deficient mice suggested that Ctr2 functions as an indirect regulator of Cu⁺-uptake and intracellular flux by stabilizing the biosynthesis of cleaved Ctr1. The cleaved Ctr1 is a transporter form which lacks metal binding Methionine- and Histidine-rich motifs and of consequence has decreased Cu⁺ and also cisplatin uptake function (Ohrvik et al., 2013). Therefore, high expression of Ctr2 seems to be associated with resistance to the cytotoxic effect of cisplatin (Blair et al., 2009) and knockdown of Ctr2 was associated with an increased cisplatin accumulation and cytotoxicity (Blair et al., 2009, 2011).


Copper-Transporting ATPase 1 and 2 (ATP7A and ATP7B)

The P-type copper-transporting ATPases ATP7A and ATP7B are also involved in cellular cisplatin handling. These transporters play an important role in regulating the cellular copper levels, because too high intracellular copper concentrations are toxic for the cell. Inactivation of these transporters, as present for example in Menkes' disease (inactivation of ATP7A) and in Wilson's disease (inactivation of ATP7B), is associated with copper deficiency because of impaired copper efflux from enterocytes into the blood or massive cellular copper overload, respectively (Gupta and Lutsenko, 2009). While ATP7A is mainly expressed in intestine, choroid plexus, vascular smooth muscle and endothelial cells, as well as in cerebrovascular endothelial cells (Lutsenko et al., 2007), ATP7B is principally expressed in the liver and the brain (Lutsenko et al., 2007). Regarding the transport of cisplatin, ATP7A and ATP7B mediate its efflux from the cell or its distribution to specific sub-cellular compartments (Katano et al., 2003, 2004; Samimi et al., 2004a,b; Safaei et al., 2008; Safaei et al., 2012). For this reason, the expression of these transporters is correlated with cisplatin cellular sensitivity and resistance (Komatsu et al., 2000; Nakagawa et al., 2008): ATP7B is stronger associated with the acquisition of resistance than Ctr1 or ATP7A (Yoshizawa et al., 2007; Mangala et al., 2009). Besides cisplatin, ATP7A and B transporters also interact with carboplatin and oxaliplatin (Samimi et al., 2004b; Martinez-Balibrea et al., 2009). Even though the effects of ATP7A and B transporters on cisplatin cellular distribution are very similar to those observed for copper, platinum drugs are not readily exported after vesicular sequestration (Samimi et al., 2004b).

Interestingly, copper transport systems are expressed and active in DRG, which are sensitive to toxicity from platinum derivatives. Here, Ctr1 is expressed in large-sized neurons and ATP7A in small DRG neurons (Ip et al., 2010), suggesting that large neurons are especially sensitive and small neurons are protected from toxic effects of platinum derivatives.

Organic Cation Transporters (OCT1-3, SLC22A1-3)

A specific interaction of cisplatin with OCTs has also been demonstrated (Ciarimboli et al., 2005; Yonezawa et al., 2005; Filipinski et al., 2008). Since OCTs have a specific organ distribution, with high renal expression, the cisplatin-OCT interaction is of special interest to explain selective organ toxicity of cisplatin. OCTs are highly expressed in excretory organs such as the liver and the kidneys, where they mediate the electrogenic uptake of their substrates in hepatocytes and proximal tubule cells, respectively (Ciarimboli, 2008). OCTs are defined as polyspecific transporters, because they can transport several unrelated substances. The driving force for the cellular transport by OCTs is the electrochemical gradient of the substrate (Ciarimboli and Schlatter, 2005). In excretory organs, OCTs mediate the first step of secretion process, consisting of substrate uptake through the basolateral plasma membrane (the blood-faced part of plasma membrane). The subsequent substrate efflux through the luminal membrane (the bile- or urine-faced part of plasma membrane) is the final secretion step, resulting in a vectorial substrate movement from the blood to the bile or urine in the liver or kidneys (**Figure 1**), respectively. In humans the paralogs hOCT1 and hOCT2 are specifically expressed in the basolateral membrane of hepatocytes and renal proximal tubule cells, respectively (Koepsell, 2004). Cisplatin seems to interact preferentially with hOCT2 (Ciarimboli et al., 2005), suggesting that hOCT2 is the critical transporter for renal cisplatin uptake in humans (**Figure 1**). Also the second- and third generation platinum-derivatives oxaliplatin (Yonezawa et al., 2006) and picoplatin (More et al., 2010), respectively, are substrates of OCTs. For the interpretation of translational studies, it is important to underline that the rodent OCT orthologs have a different organ distribution and kinetic properties compared with human OCTs: for example, in mice OCT1 is expressed in renal proximal tubules at higher level than OCT2 (Schlatter et al., 2014). Competition of OCT-mediated cisplatin transport is able to reduce cisplatin uptake (Ciarimboli et al., 2010; Wehe et al., 2014) and toxicity (Ciarimboli et al., 2005, 2010; Yonezawa et al., 2005) *in vitro* and *in vivo*. OCT2 has been demonstrated to be expressed in the mouse cochlea in hair cells of Corti organ and in the cells of the stria vascularis (Ciarimboli et al., 2010) and in mouse and human DRG (Sprowl et al., 2013), structures that are specially sensitive to toxicity by platinum-derivatives. In animal models it has been demonstrated that OCTs are critical mediators of cisplatin ototoxicity (Ciarimboli et al., 2010) and oxaliplatin peripheral neurotoxicity (Sprowl et al., 2013). Currently, the role of genetic variations of SLC22A2 for the development of cisplatin-induced nephrotoxicity is debated: some publications describe an association of the SLC22A2 non-synonymous single nucleotide polymorphism (SNP) 808 G/T (resulting in an exchange of alanine to serine at position 270) with a protection against cisplatin nephrotoxicity (Filipinski et al., 2009; Zhang and Zhou, 2012), other studies however could not confirm this finding (Tzvetkov et al., 2011). Recent results suggest that the SNP 808 G/T confers a protection against cisplatin ototoxicity in a pediatric population (Lanvers-Kaminsky et al., 2015). Therefore,



it could be assumed that inhibition of hOCT2 by a competitor could prevent cisplatin-induced side effects. Indeed, there are several indications that patient co-treatment with cimetidine protects the kidneys from cisplatin nephrotoxicity (Sleijfer et al., 1987; Zhang and Zhou, 2012). However, such a protective therapeutic option is acceptable, if the uptake of cisplatin into target tumor cells is not compromised. In animal models and in patients, it has been demonstrated that the expression of OCTs in tumor cells is down-regulated by epigenetic modifications (Schaeffeler et al., 2011; Chen et al., 2013; Yang et al., 2013). Moreover, studies in an animal model have shown that co-treatment with cimetidine did not change the antitumor efficacy of cisplatin (Katsuda et al., 2010). For these reasons, inhibition of OCTs remains an attractive option to protect patients from cisplatin side effects.

Comparing the transport properties of OCTs and Ctr1, it was underlined that Ctr1 is an equilibrative transporter. On the other side, although OCTs are not active transporters in the sense of their activity being directly dependent on ATP (primary active) or a trans-membrane ion flux (secondary active), the plasma membrane potential creates both an electrical and a concentration gradient for their cationic substrates. This allows an accumulation of platinum derivatives in the cells to intracellular concentrations exceeding the extracellular level (Yonezawa and Inui, 2011). However, the affinities of OCT2 and Ctr1 for cisplatin seem to be similar [11 (Filipski et al., 2008) and 17 μ M (Liang et al., 2009), respectively].

While OCTs mediate the basolateral uptake of cisplatin in renal proximal tubule cells, other transporters expressed in the apical cell membrane are involved in the cisplatin secretion into the urine (Figure 1).

Multidrug and Toxin Extrusion Protein 1 (MATE1, SLC47A1)

Several evidences indicate that MATE1 mediates secretion of cisplatin into the urine. Mice with genetic deletion of MATE1 are more sensitive to cisplatin nephrotoxicity (Nakamura et al., 2010). Furthermore, cell transfected with MATE1 displayed a higher cisplatin uptake than control cells (Nakamura et al., 2010). Interestingly, MATE1 and MATE2-K, another member of MATE family which is solely expressed in human kidneys, seem to transport oxaliplatin with higher affinity than cisplatin (Yokoo et al., 2007), offering a possible explanation of the low oxaliplatin nephrotoxicity. As outlined above, inhibition of OCT2 may be a protective strategy against cisplatin nephrotoxicity. However, some inhibitors of OCT2 such as cimetidine (Matsushima et al., 2009) and ondansetron (Li et al., 2013) interact with higher potency with MATE1, blocking cisplatin efflux from the cells and potentially increasing cisplatin renal toxicity. Indeed, co-treatment of mice with cisplatin and cimetidine was effective in protecting the animals from ototoxicity but not from nephrotoxicity (Ciarimboli et al., 2010).

There are some investigations suggesting a role of novel organic cation transporters (OCTNs) for oxaliplatin transport. These transporters are expressed on the apical membrane of renal proximal tubule cells (Ciarimboli, 2008) and in rat DRGs (Jong et al., 2011). When transfected in human embryonic kidney cells, rat and human OCTN1 and OCTN2 mediate significant oxaliplatin uptake (Jong et al., 2011), suggesting that OCTNs are involved in oxaliplatin neurotoxicity. Apart from these not directly ATP-dependent transporters, multidrug resistance-associated protein 2 (Mrp2) transporter

seems to be involved in the efflux of cisplatin and its conjugates from kidney cells, and for this reason to play an important role for control of cisplatin renal toxicity (Wen et al., 2014).

In conclusion, cellular transport of platinum derivatives is mediated by several transport systems. Some transporters, such as OCTs, are specifically expressed in organs, which are damaged by antitumor therapy with platinum derivatives. For this reason, they may be a target for protective intervention. However,

an efficient protection can be only reached by specific inhibition of OCTs.

Acknowledgments

The support of the “Interdisziplinäres Zentrum für Klinische Forschung (IZKF)” (CIA02/013/13) at the Medical Faculty of Münster University for the studies of the author on OCT interactions with drugs is acknowledged.

References

- Adelsberger, H., Quasthoff, S., Grosskreutz, J., Lepier, A., Eckel, F., and Lersch, C. (2000). The chemotherapeutic oxaliplatin alters voltage-gated Na(+) channel kinetics on rat sensory neurons. *Eur. J. Pharmacol.* 406, 25–32. doi: 10.1016/S0014-2999(00)00667-1
- Albers, J. W., Chaudhry, V., Cavaletti, G., and Donehower, R. C. (2011). Interventions for preventing neuropathy caused by cisplatin and related compounds. *Cochrane Database Syst. Rev.* 2:CD005228. doi: 10.1002/14651858.CD005228.pub3
- Amptoulach, S., and Tsavaris, N. (2011). Neurotoxicity caused by the treatment with platinum analogues. *Chemother. Res. Pract.* 2011:843019. doi: 10.1155/2011/843019
- Blair, B. G., Larson, C. A., Adams, P. L., Abada, P. B., Pesce, C. E., Safaei, R., et al. (2011). Copper transporter 2 regulates endocytosis and controls tumor growth and sensitivity to cisplatin *in vivo*. *Mol. Pharmacol.* 79, 157–166. doi: 10.1124/mol.110.068411
- Blair, B. G., Larson, C. A., Safaei, R., and Howell, S. B. (2009). Copper transporter 2 regulates the cellular accumulation and cytotoxicity of Cisplatin and Carboplatin. *Clin. Cancer Res.* 15, 4312–4321. doi: 10.1158/1078-0432.CCR-09-0311
- Bregman, C. L., and Williams, P. D. (1986). Comparative nephrotoxicity of carboplatin and cisplatin in combination with tobramycin. *Cancer Chemother. Pharmacol.* 18, 117–123. doi: 10.1007/BF00262279
- Cano, J. R., Catalan, B., and Jara, C. (1998). [Neuropathy due to cisplatin]. *Rev. Neurol.* 27, 606–610.
- Carozzi, V. A., Marmiroli, P., and Cavaletti, G. (2010). The role of oxidative stress and anti-oxidant treatment in platinum-induced peripheral neurotoxicity. *Curr. Cancer Drug Targets* 10, 670–682. doi: 10.2174/156800910793605820
- Cassidy, J., and Misset, J. L. (2002). Oxaliplatin-related side effects: characteristics and management. *Semin. Oncol.* 29, 11–20. doi: 10.1053/sonc.2002.35524
- Cavaletti, G., and Zanna, C. (2002). Current status and future prospects for the treatment of chemotherapy-induced peripheral neurotoxicity. *Eur. J. Cancer* 38, 1832–1837. doi: 10.1016/S0959-8049(02)00229-0
- Chen, L., Hong, C., Chen, E. C., Yee, S. W., Xu, L., Almof, E. U., et al. (2013). Genetic and epigenetic regulation of the organic cation transporter 3, SLC22A3. *Pharmacogenomics J.* 13, 110–120. doi: 10.1038/tpj.2011.60
- Ciarimboli, G. (2008). Organic cation transporters. *Xenobiotica* 38, 936–971. doi: 10.1080/00498250701882482
- Ciarimboli, G. (2011). Role of organic cation transporters in drug-induced toxicity. *Expert Opin. Drug Metab. Toxicol.* 7, 159–174. doi: 10.1517/17425255.2011.547474
- Ciarimboli, G., Deuster, D., Knief, A., Sperling, M., Holtkamp, M., Edemir, B., et al. (2010). Organic cation transporter 2 mediates cisplatin-induced oto- and nephrotoxicity and is a target for protective interventions. *Am. J. Pathol.* 176, 1169–1180. doi: 10.2353/ajpath.2010.090610
- Ciarimboli, G., Ludwig, T., Lang, D., Pavenstädt, H., Koepsell, H., Piechota, H. J., et al. (2005). Cisplatin nephrotoxicity is critically mediated via the human organic cation transporter 2. *Am. J. Pathol.* 167, 1477–1484. doi: 10.1016/S0002-9440(10)61234-5
- Ciarimboli, G., and Schlatter, E. (2005). Regulation of organic cation transport. *Pflugers Arch.* 449, 423–441. doi: 10.1007/s00424-004-1355-5
- Daugaard, G. (1990). Cisplatin nephrotoxicity: experimental and clinical studies. *Dan. Med. Bull.* 37, 1–12.
- Dobyan, D. C., Levi, J., Jacobs, C., Kosek, J., and Weiner, M. W. (1980). Mechanism of cis-platinum nephrotoxicity: II. Morphologic observations. *J. Pharmacol. Exp. Ther.* 213, 551–556.
- Einhorn, L. H. (2002). Curing metastatic testicular cancer. *Proc. Natl. Acad. Sci. U.S.A.* 99, 4592–4595. doi: 10.1073/pnas.072067999
- Filipski, K. K., Loos, W. J., Verweij, J., and Sparreboom, A. (2008). Interaction of Cisplatin with the human organic cation transporter 2. *Clin. Cancer Res.* 14, 3875–3880. doi: 10.1158/1078-0432.CCR-07-4793
- Filipski, K. K., Mathijssen, R. H., Mikkelsen, T. S., Schinkel, A. H., and Sparreboom, A. (2009). Contribution of organic cation transporter 2 (OCT2) to cisplatin-induced nephrotoxicity. *Clin. Pharmacol. Ther.* 86, 396–402. doi: 10.1038/clpt.2009.139
- Gamelin, E., Gamelin, L., Bossi, L., and Quasthoff, S. (2002). Clinical aspects and molecular basis of oxaliplatin neurotoxicity: current management and development of preventive measures. *Semin. Oncol.* 29, 21–33. doi: 10.1053/sonc.2002.35525
- Goren, M. P., Wright, R. K., and Horowitz, M. E. (1986). Cumulative renal tubular damage associated with cisplatin nephrotoxicity. *Cancer Chemother. Pharmacol.* 18, 69–73. doi: 10.1007/BF00253068
- Gupta, A., and Lutsenko, S. (2009). Human copper transporters: mechanism, role in human diseases and therapeutic potential. *Future Med. Chem.* 1, 1125–1142. doi: 10.4155/fmc.09.84
- Hartmann, J. T., and Lipp, H. P. (2003). Toxicity of platinum compounds. *Expert Opin. Pharmacother.* 4, 889–901. doi: 10.1517/14656566.4.6.889
- Hellberg, V., Wallin, I., Eriksson, S., Hernlund, E., Jerremalm, E., Berndtsson, M., et al. (2009). Cisplatin and oxaliplatin toxicity: importance of cochlear kinetics as a determinant for ototoxicity. *J. Natl. Cancer Inst.* 101, 37–47. doi: 10.1093/jnci/djn418
- Holzer, A. K., Manorek, G. H., and Howell, S. B. (2006). Contribution of the major copper influx transporter CTR1 to the cellular accumulation of cisplatin, carboplatin, and oxaliplatin. *Mol. Pharmacol.* 70, 1390–1394. doi: 10.1124/mol.106.022624
- Holzer, A. K., Samimi, G., Katano, K., Naerdemann, W., Lin, X., Safaei, R., et al. (2004). The copper influx transporter human copper transport protein 1 regulates the uptake of cisplatin in human ovarian carcinoma cells. *Mol. Pharmacol.* 66, 817–823. doi: 10.1124/mol.104.001198
- Houjou, T., Nakano, K., Ike, O., Wada, H., Hitomi, S., Shinmi, Y., et al. (1996). Oral sustained-release cisplatin capsule. *J. Pharm. Pharmacol.* 48, 474–478. doi: 10.1111/j.2042-7158.1996.tb05957.x
- Ip, V., Liu, J. J., Mercer, J. F., and McKeage, M. J. (2010). Differential expression of ATP7A, ATP7B and CTR1 in adult rat dorsal root ganglion tissue. *Mol. Pain* 6, 53. doi: 10.1186/1744-8069-6-53
- Ishida, S., Lee, J., Thiele, D. J., and Herskowitz, I. (2002). Uptake of the anticancer drug cisplatin mediated by the copper transporter Ctr1 in yeast and mammals. *Proc. Natl. Acad. Sci. U.S.A.* 99, 14298–14302. doi: 10.1073/pnas.162491399
- Jacobs, C., Coleman, C. N., Rich, L., Hirst, K., and Weiner, M. W. (1984). Inhibition of cis-diamminedichloroplatinum secretion by the human kidney with probenecid. *Cancer Res.* 44, 3632–3635.
- Jong, N. N., Nakanishi, T., Liu, J. J., Tamai, I., and McKeage, M. J. (2011). Oxaliplatin transport mediated by organic cation/carnitine transporters OCTN1 and OCTN2 in overexpressing human embryonic kidney 293 cells and rat dorsal root ganglion neurons. *J. Pharmacol. Exp. Ther.* 338, 537–547. doi: 10.1124/jpet.111.181297

- Katano, K., Safaei, R., Samimi, G., Holzer, A., Rochdi, M., and Howell, S. B. (2003). The copper export pump ATP7B modulates the cellular pharmacology of carboplatin in ovarian carcinoma cells. *Mol. Pharmacol.* 64, 466–473. doi: 10.1124/mol.64.2.466
- Katano, K., Safaei, R., Samimi, G., Holzer, A., Tomioka, M., Goodman, M., et al. (2004). Confocal microscopic analysis of the interaction between cisplatin and the copper transporter ATP7B in human ovarian carcinoma cells. *Clin. Cancer Res.* 10, 4578–4588. doi: 10.1158/1078-0432.CCR-03-0689
- Katsuda, H., Yamashita, M., Katsura, H., Yu, J., Waki, Y., Nagata, N., et al. (2010). Protecting Cisplatin-induced nephrotoxicity with cimetidine does not affect antitumor activity. *Biol. Pharm. Bull.* 33, 1867–1871. doi: 10.1248/bpb.33.1867
- Koepsell, H. (2004). Polyspecific organic cation transporters: their functions and interactions with drugs. *Trends Pharmacol. Sci.* 25, 375–381. doi: 10.1016/j.tips.2004.05.005
- Komatsu, M., Sumizawa, T., Mutoh, M., Chen, Z. S., Terada, K., Furukawa, T., et al. (2000). Copper-transporting P-type adenosine triphosphatase (ATP7B) is associated with cisplatin resistance. *Cancer Res.* 60, 1312–1316.
- Lanvers-Kaminsky, C., Sprowl, J. A., Malath, I., Deuster, D., Eveslage, M., Schlatter, E., et al. (2015). Human organic cation transporter 2 variant c.808G>T carriers are protected against Cisplatin-induced ototoxicity. *Pharmacogenomics* 16, 323–332. doi: 10.2217/pgs.14.182
- Larson, C. A., Blair, B. G., Safaei, R., and Howell, S. B. (2009). The role of the mammalian copper transporter 1 in the cellular accumulation of platinum-based drugs. *Mol. Pharmacol.* 75, 324–330. doi: 10.1124/mol.108.052381
- Lee, Y. Y., Choi, C. H., Do, I. G., Song, S. Y., Lee, W., Park, H. S., et al. (2011). Prognostic value of the copper transporters, CTR1 and CTR2, in patients with ovarian carcinoma receiving platinum-based chemotherapy. *Gynecol. Oncol.* 122, 361–365. doi: 10.1016/j.ygyno.2011.04.025
- Li, Q., Guo, D., Dong, Z., Zhang, W., Zhang, L., Huang, S. M., et al. (2013). Ondansetron can enhance cisplatin-induced nephrotoxicity via inhibition of multiple toxin and extrusion proteins (MATEs). *Toxicol. Appl. Pharmacol.* 273, 100–109. doi: 10.1016/j.taap.2013.08.024
- Liang, Z. D., Long, Y., Tsai, W. B., Fu, S., Kurzrock, R., Gagea-Iurascu, M., et al. (2012). Mechanistic basis for overcoming platinum resistance using copper chelating agents. *Mol. Cancer Ther.* 11, 2483–2494. doi: 10.1158/1535-7163.MCT-12-0580
- Liang, Z. D., Stockton, D., Savaraj, N., and Tien, K. M. (2009). Mechanistic comparison of human high-affinity copper transporter 1-mediated transport between copper ion and cisplatin. *Mol. Pharmacol.* 76, 843–853. doi: 10.1124/mol.109.056416
- Lin, X., Okuda, T., Holzer, A., and Howell, S. B. (2002). The copper transporter CTR1 regulates cisplatin uptake in *Saccharomyces cerevisiae*. *Mol. Pharmacol.* 62, 1154–1159. doi: 10.1124/mol.62.5.1154
- Liu, J. J., Jamieson, S. M., Subramaniam, J., Ip, V., Jong, N. N., Mercer, J. F., et al. (2009). Neuronal expression of copper transporter 1 in rat dorsal root ganglia: association with platinum neurotoxicity. *Cancer Chemother. Pharmacol.* 64, 847–856. doi: 10.1007/s00280-009-1017-6
- Lutsenko, S., Barnes, N. L., Bartee, M. Y., and Dmitriev, O. Y. (2007). Function and regulation of human copper-transporting ATPases. *Physiol. Rev.* 87, 1011–1046. doi: 10.1152/physrev.00004.2006
- Mangala, L. S., Zuzel, V., Schmandt, R., Leshane, E. S., Halder, J. B., Armaiz-Pena, G. N., et al. (2009). Therapeutic targeting of ATP7B in ovarian carcinoma. *Clin. Cancer Res.* 15, 3770–3780. doi: 10.1158/1078-0432.CCR-08-2306
- Martinez-Balibrea, E., Martinez-Cardus, A., Musulen, E., Gines, A., Manzano, J. L., Aranda, E., et al. (2009). Increased levels of copper efflux transporter ATP7B are associated with poor outcome in colorectal cancer patients receiving oxaliplatin-based chemotherapy. *Int. J. Cancer* 124, 2905–2910. doi: 10.1002/ijc.24273
- Matsushima, S., Maeda, K., Inoue, K., Ohta, K. Y., Yuasa, H., Kondo, T., et al. (2009). The inhibition of human multidrug and toxin extrusion 1 is involved in the drug-drug interaction caused by cimetidine. *Drug Metab. Dispos.* 37, 555–559. doi: 10.1124/dmd.108.023911
- Meijer, C., de Vries, E. G., Marmioli, P., Tredici, G., Frattola, L., and Cavaletti, G. (1999). Cisplatin-induced DNA-platination in experimental dorsal root ganglia neuronopathy. *Neurotoxicology* 20, 883–887.
- More, S. S., Li, S., Yee, S. W., Chen, L., Xu, Z., Jablons, D. M., et al. (2010). Organic cation transporters modulate the uptake and cytotoxicity of picoplatin, a third-generation platinum analogue. *Mol. Cancer Ther.* 9, 1058–1069. doi: 10.1158/1535-7163.MCT-09-1084
- Musial-Bright, L., Fengler, R., Henze, G., and Hernaiz, D. P. (2011). Carboplatin and ototoxicity: hearing loss rates among survivors of childhood medulloblastoma. *Childs Nerv. Syst.* 27, 407–413. doi: 10.1007/s00381-010-1300-1
- Nakagawa, T., Inoue, Y., Kodama, H., Yamazaki, H., Kawai, K., Suemizu, H., et al. (2008). Expression of copper-transporting P-type adenosine triphosphatase (ATP7B) correlates with cisplatin resistance in human non-small cell lung cancer xenografts. *Oncol. Rep.* 20, 265–270. doi: 10.3892/or_00000002
- Nakamura, T., Yonezawa, A., Hashimoto, S., Katsura, T., and Inui, K. I. (2010). Disruption of multidrug and toxin extrusion MATE1 potentiates cisplatin-induced nephrotoxicity. *Biochem. Pharmacol.* 80, 1762–1767. doi: 10.1016/j.bcp.2010.08.019
- Ohrvik, H., Nose, Y., Wood, L. K., Kim, B. E., Gleber, S. C., Ralle, M., et al. (2013). Ctr2 regulates biogenesis of a cleaved form of mammalian Ctr1 metal transporter lacking the copper- and cisplatin-binding ecto-domain. *Proc. Natl. Acad. Sci. U.S.A.* 110, E4279–E4288. doi: 10.1073/pnas.1311749110
- Ohrvik, H., and Thiele, D. J. (2014). The role of Ctr1 and Ctr2 in mammalian copper homeostasis and platinum-based chemotherapy. *J. Trace Elem. Med. Biol.* doi: 10.1016/j.jtemb.2014.03.006. [Epub ahead of print].
- Pabla, N., and Dong, Z. (2008). Cisplatin nephrotoxicity: mechanisms and renoprotective strategies. *Kidney Int.* 73, 994–1007. doi: 10.1038/sj.ki.5002786
- Price, P. M., Safirstein, R. L., and Megyesi, J. (2004). Protection of renal cells from cisplatin toxicity by cell cycle inhibitors. *Am. J. Physiol. Renal Physiol.* 286, F378–F384. doi: 10.1152/ajprenal.00192.2003
- Rabik, C. A., and Dolan, M. E. (2007). Molecular mechanisms of resistance and toxicity associated with platinating agents. *Cancer Treat. Rev.* 33, 9–23. doi: 10.1016/j.ctrv.2006.09.006
- Reddel, R. R., Kefford, R. F., Grant, J. M., Coates, A. S., Fox, R. M., and Tattersall, M. H. (1982). Ototoxicity in patients receiving cisplatin: importance of dose and method of drug administration. *Cancer Treat. Rep.* 66, 19–23.
- Renes, J., de Vries, E. G., Jansen, P. L., and Muller, M. (2000). The (patho)physiological functions of the MRP family. *Drug Resist. Updat.* 3, 289–302. doi: 10.1054/drup.2000.0156
- Roelofs, R. I., Hrushesky, W., Rogin, J., and Rosenberg, L. (1984). Peripheral sensory neuropathy and cisplatin chemotherapy. *Neurology* 34, 934–938. doi: 10.1212/WNL.34.7.934
- Rybik, L. P., and Ramkumar, V. (2007). Ototoxicity. *Kidney Int.* 72, 931–935. doi: 10.1038/sj.ki.5002434
- Safaei, R., Adams, P. L., Maktabi, M. H., Mathews, R. A., and Howell, S. B. (2012). The CXXC motifs in the metal binding domains are required for ATP7B to mediate resistance to cisplatin. *J. Inorg. Biochem.* 110, 8–17. doi: 10.1016/j.jinorgbio.2012.02.016
- Safaei, R., and Howell, S. B. (2005). Copper transporters regulate the cellular pharmacology and sensitivity to Pt drugs. *Crit. Rev. Oncol. Hematol.* 53, 13–23. doi: 10.1016/j.critrevonc.2004.09.007
- Safaei, R., Otani, S., Larson, B. J., Rasmussen, M. L., and Howell, S. B. (2008). Transport of cisplatin by the copper efflux transporter ATP7B. *Mol. Pharmacol.* 73, 461–468. doi: 10.1124/mol.107.040980
- Samimi, G., Katano, K., Holzer, A. K., Safaei, R., and Howell, S. B. (2004b). Modulation of the cellular pharmacology of cisplatin and its analogs by the copper exporters ATP7A and ATP7B. *Mol. Pharmacol.* 66, 25–32. doi: 10.1124/mol.66.1.25
- Samimi, G., Safaei, R., Katano, K., Holzer, A. K., Rochdi, M., Tomioka, M., et al. (2004a). Increased expression of the copper efflux transporter ATP7A mediates resistance to cisplatin, carboplatin, and oxaliplatin in ovarian cancer cells. *Clin. Cancer Res.* 10, 4661–4669. doi: 10.1158/1078-0432.CCR-04-0137
- Schaeffeler, E., Hellerbrand, C., Nies, A. T., Winter, S., Kruck, S., Hofmann, U., et al. (2011). DNA methylation is associated with downregulation of the organic cation transporter OCT1 (SLC22A1) in human hepatocellular carcinoma. *Genome Med.* 3, 82. doi: 10.1186/gm298
- Schlatter, E., Klassen, P., Massmann, V., Holle, S. K., Guckel, D., Edemir, B., et al. (2014). Mouse organic cation transporter 1 determines properties and regulation of basolateral organic cation transport in renal proximal tubules. *Pflügers Arch.* 466, 1581–1589. doi: 10.1007/s00424-013-1395-9
- Scrceni, D., and McKeage, M. J. (1999). Platinum neurotoxicity: clinical profiles, experimental models and neuroprotective approaches. *J. Inorg. Biochem.* 77, 105–110. doi: 10.1016/S0162-0134(99)00135-X

- Sinani, D., Adle, D. J., Kim, H., and Lee, J. (2007). Distinct mechanisms for Ctr1-mediated copper and cisplatin transport. *J. Biol. Chem.* 282, 26775–26785. doi: 10.1074/jbc.M703973200
- Skinner, R. (2004). Best practice in assessing ototoxicity in children with cancer. *Eur. J. Cancer* 40, 2352–2354. doi: 10.1016/j.ejca.2004.08.002
- Sleijfer, D. T., Offerman, J. J., Mulder, N. H., Verweij, M., van der Hem, G. K., Schraffordt Koops, H. S., et al. (1987). The protective potential of the combination of verapamil and cimetidine on cisplatin-induced nephrotoxicity in man. *Cancer* 60, 2823–2828.
- Sprowl, J. A., Ciarimboli, G., Lancaster, C. S., Giovinazzo, H., Gibson, A. A., Du, G., et al. (2013). Oxaliplatin-induced neurotoxicity is dependent on the organic cation transporter OCT2. *Proc. Natl. Acad. Sci. U.S.A.* 110, 11199–11204. doi: 10.1073/pnas.1305321110
- Takeshita, M., Banno, Y., Nakamura, M., Otsuka, M., Teramachi, H., Tsuchiya, T., et al. (2011). The pivotal role of intracellular calcium in oxaliplatin-induced inhibition of neurite outgrowth but not cell death in differentiated PC12 cells. *Chem. Res. Toxicol.* 24, 1845–1852. doi: 10.1021/tx200160g
- Thompson, S. W., Davis, L. E., Kornfeld, M., Hilgers, R. D., and Standefer, J. C. (1984). Cisplatin neuropathy. Clinical, electrophysiologic, morphologic, and toxicologic studies. *Cancer* 54, 1269–1275.
- Tsai, C. Y., Larson, C. A., Safaei, R., and Howell, S. B. (2014). Molecular modulation of the copper and cisplatin transport function of CTR1 and its interaction with IRS-4. *Biochem. Pharmacol.* 90, 379–387. doi: 10.1016/j.bcp.2014.06.019
- Tzvetkov, M. V., Behrens, G., O'Brien, V. P., Hohloch, K., Brockmoller, J., and Benohr, P. (2011). Pharmacogenetic analyses of cisplatin-induced nephrotoxicity indicate a renoprotective effect of ERCC1 polymorphisms. *Pharmacogenomics* 12, 1417–1427. doi: 10.2217/pgs.11.93
- van den Berghe, P. V., Folmer, D. E., Malingre, H. E., van Beurden, E., and Klomp, A. E., van de Sluis, B., et al. (2007). Human copper transporter 2 is localized in late endosomes and lysosomes and facilitates cellular copper uptake. *Biochem. J.* 407, 49–59. doi: 10.1042/BJ20070705
- Wang, D., and Lippard, S. J. (2005). Cellular processing of platinum anticancer drugs. *Nat. Rev. Drug Discov.* 4, 307–320. doi: 10.1038/nrd1691
- Wehe, C. A., Beyer, G., Sperling, M., Ciarimboli, G., and Karst, U. (2014). Assessing the intracellular concentration of platinum in medulloblastoma cell lines after Cisplatin incubation. *J. Trace Elem. Med. Biol.* 28, 166–172. doi: 10.1016/j.jtemb.2014.01.001
- Wen, X., Buckley, B., McCandlish, E., Goedken, M. J., Syed, S., Pelis, R., et al. (2014). Transgenic expression of the human MRP2 transporter reduces cisplatin accumulation and nephrotoxicity in Mrp2-null mice. *Am. J. Pathol.* 184, 1299–1308. doi: 10.1016/j.ajpath.2014.01.025
- Xu, X., Duan, L., Zhou, B., Ma, R., Zhou, H., and Liu, Z. Q. (2012). Genetic polymorphism of copper transporter protein 1 is related to platinum resistance in Chinese non-small cell lung carcinoma patients. *Clin. Exp. Pharmacol. Physiol.* 39, 786–792. doi: 10.1111/j.1440-1681.2012.05741.x
- Yang, J., Kalogerou, M., Gallacher, J., Sampson, J. R., and Shen, M. H. (2013). Renal tumours in a Tsc1+/- mouse model show epigenetic suppression of organic cation transporters Slc22a1, Slc22a2 and Slc22a3, and do not respond to metformin. *Eur. J. Cancer* 49, 1479–1490. doi: 10.1016/j.ejca.2012.10.027
- Yasumasu, T., Ueda, T., Uozumi, J., Mihara, Y., and Kumazawa, J. (1992). Comparative study of cisplatin and carboplatin on pharmacokinetics, nephrotoxicity and effect on renal nuclear DNA synthesis in rats. *Pharmacol. Toxicol.* 70, 143–147. doi: 10.1111/j.1600-0773.1992.tb00445.x
- Yokoo, S., Yonezawa, A., Masuda, S., Fukatsu, A., Katsura, T., and Inui, K. (2007). Differential contribution of organic cation transporters, OCT2 and MATE1, in platinum agent-induced nephrotoxicity. *Biochem. Pharmacol.* 74, 477–487. doi: 10.1016/j.bcp.2007.03.004
- Yonezawa, A., and Inui, K. (2011). Organic cation transporter OCT/SLC22A and H(+)/organic cation antiporter MATE/SLC47A are key molecules for nephrotoxicity of platinum agents. *Biochem. Pharmacol.* 81, 563–568. doi: 10.1016/j.bcp.2010.11.016
- Yonezawa, A., Masuda, S., Nishihara, K., Yano, I., Katsura, T., and Inui, K. (2005). Association between tubular toxicity of cisplatin and expression of organic cation transporter rOCT2 (Slc22a2) in the rat. *Biochem. Pharmacol.* 70, 1823–1831. doi: 10.1016/j.bcp.2005.09.020
- Yonezawa, A., Masuda, S., Yokoo, S., Katsura, T., and Inui, K. (2006). Cisplatin and oxaliplatin, but not carboplatin and nedaplatin, are substrates for human organic cation transporters (SLC22A1-3 and multidrug and toxin extrusion family). *J. Pharmacol. Exp. Ther.* 319, 879–886. doi: 10.1124/jpet.106.110346
- Yoshizawa, K., Nozaki, S., Kitahara, H., Ohara, T., Kato, K., Kawashiri, S., et al. (2007). Copper efflux transporter (ATP7B) contributes to the acquisition of cisplatin-resistance in human oral squamous cell lines. *Oncol. Rep.* 18, 987–991. doi: 10.3892/or.18.4.987
- Zhang, J., and Zhou, W. (2012). Ameliorative effects of SLC22A2 gene polymorphism 808 G/T and cimetidine on cisplatin-induced nephrotoxicity in Chinese cancer patients. *Food Chem. Toxicol.* 50, 2289–2293. doi: 10.1016/j.fct.2012.03.077

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Harrach and Ciarimboli. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.